

# WATER ECOSYSTEMS

## The slow death of Lake Sibaya

*Why is the water level dropping steadily in one of South Africa's most valuable freshwater lakes? And what can be done to stop it from drying up further? Article by Tony Carnie.*



Lake Sibaya, named after the Zulu word for **cattle kraal**, is the country's largest coastal freshwater lake.

With a surface area of more than 60 km<sup>2</sup> and a water depth of 41 m at the deepest point, it has been estimated that Sibaya can hold more water than all three large dams in the uMngeni River system (Midmar, Albert Falls and Inanda dams) or about 44% of the Pongolapoort Dam.

The lake was once an ancient river estuary, before it was cut off from the Indian Ocean by the gradual formation of a coastal dune barrier. Located in the northern part of KwaZulu-Natal, about 60 km south of the Mozambique border, Sibaya lies on a flat coastal plain where most of the surrounding landscape rarely

rises more than 100 m above sea level.

But beneath the region's largely infertile and sandy soils there is a vital groundwater aquifer which has helped to sustain local communities and the ecology around Lake Sibaya for millennia. As part of this aquifer, the lake is an almost direct reflection of the regionally-important groundwater table.

Now the lake is in peril, drying out rapidly due to drought, increasing human water demands and the thirsty roots of exotic timber plantations that have been dubbed "money trees".

During the severe 2014-2016 drought, Lake Sibaya's water level dropped quickly – along with that of several other lakes and

dams in the region. But whereas Lake St Lucia and local dams have improved gradually since the drought, Sibaya has not risen – and is still dropping.

According to Van Rensburg, the level of the lake reached the lowest recorded level in 2014 and has continued to decline since then. Van Rensburg, the coordinator of the Grasslands-Wetlands-Forests Node of the SA Environmental Observation Network (SAEON), says the main section of the lake has dropped over five meters over the last decade when historic fluctuations were only between one and three meters. Because of the surface area of the lake, even a small drop in water level equates to a large volume of water lost.

Having separated from the main lake for the first time in living memory in 2015, the southern basin – which supplies Mbazwana and Sodwana Bay – has now dropped even lower than the main lake.

One of the main reasons for this, explains Van Rensburg, is because Sibaya is almost entirely dependent on local rainfall and groundwater recharge. Unlike open river estuaries (which receive both seawater inflow, and freshwater from surrounding rivers) Sibaya has no large perennial rivers that import water into the system from upper catchments. If it does not rain locally, there is no recharge to the system.


Located in quaternary catchment W70A, the total catchment

area for the lake is estimated at 530 km<sup>2</sup>, of which 60-70 km<sup>2</sup> is taken up by the lake itself. The surrounding human population has also increased significantly in recent decades, although recent studies by Prof Jeff Smithers of the University of KwaZulu-Natal suggest that water abstractions for human use are considered to be modest.

In a study published in 2017, Smithers et al. suggested that (other than drought) a major cause of Sibaya's recent water loss can be linked to depletion by water-guzzling gum and pine plantations.

The first State forestry plantations in the Sibaya area were established in the 1950s and now cover at least 23% of the catchment. Studies show that between 2001 and 2014, the level of Lake Sibaya dropped from close to 20 m above sea level to nearly 16 m above sea level – its lowest level since the commencement of record keeping more than 50 years ago and simulations going back to 1914.

Smithers and his colleagues also modelled the hydrology back to 1986. The results indicated that approximately 35% of the drop in lake levels since 2001 could be attributed to the impact of afforestation, although they noted that there was some uncertainty regarding the exact history, extent and impact of afforestation in the catchment. In contrast, the results indicated that the impact of domestic abstractions on lake levels had been 'negligible'.



Jannie Weitz

The first State forestry plantations in the Sibaya area were established in the 1950s and now cover at least 23% of the catchment. Hydrologists have suggested that at least 5km<sup>2</sup> of plantations be removed to protect the lake's water resources.



Sue van Rensburg



*University of KwaZulu-Natal PhD student, Mlu Shabalala, conducting groundwater use management studies on early growth macadamia and eucalyptus plantations in relation to grasslands.*

“The major cause of the drop in the level of Lake Sibaya since 2001 is postulated to be the 10-year period of significantly lower than average rainfall which lasted from 2001 to 2011,” said Smithers. Subsequent analysis (Blamey et al) have shown that 2015/2016 was the driest year on record for the region.

A yield analysis demonstrated that at 2014 levels of water use from timber plantations and domestic abstractions, no sustainable additional yield was possible. The 2015/16 drought and subsequent erratic rainfall has exacerbated this situation.

Smithers suggested that at least 5 km<sup>2</sup> of forestry should be removed to provide for additional domestic abstraction. Several other studies in the region have also highlighted the significant impact of timber plantations on local water resources.

In 2018, Prof Molla Demlie of the University of KwaZulu-Natal (UKZN) reported that the large primary aquifer on South Africa’s north-eastern coastal plain was under stress from domestic water abstraction, irrigation, commercial forests and reduced rainfall linked to climate change. Prof Demlie suggested that removing water-intensive commercial plantations from the recharge area of the catchments and halting over abstraction of the freshwater lakes were some of the solutions needed to avert serious environmental impacts.

Further studies by hydrogeologist Claudia Brites in the Nyalazi plantation near Lake St Lucia suggest that deep-rooted gum trees use more than twice as much groundwater compared to

locally indigenous trees, while hydrologist Brian Rawlins reported that gum and pine plantations around St Lucia could reduce the lake’s groundwater inflow by as much as 30% during periods of extreme drought.

More recent modelling studies by UKZN hydrologist, Jannie Weitz, warned that further severe depletion of groundwater around Sibaya could lead to salt-water intruding into the lake from the Indian Ocean.

SAEON’s Sue van Rensburg has also voiced concern that climate change may be exacerbating the problem due to reduced rainfall in the catchment in recent years. Van Rensburg started visiting Sibaya regularly from 2014, capturing a set of time-series images which graphically illustrate the declining water level and gradual separation of the southern basin from the main lake.

Van Rensburg, a former regional ecologist for the Hluhluwe-Imfolozi Park who also spent two years working with communities around Serengeti National Park in Tanzania, is determined to help find a sustainable solution before it is too late: “Our job is to do science with and for society to ensure a more sustainable future”

She sees the emerging water crisis as an important research opportunity to understand the relative impacts of land-use, water abstraction, weather, climate change and sea-level change on one of the country’s most important coastal aquifer systems.



Sue van Rensburg

*SAEON technician, Siphwe Mfeka, downloads groundwater data from a monitoring sensor. Working in collaboration with various stakeholders, SAEON has established a network of groundwater monitoring sites which it maintains to determine trends in ground water dynamics and how this links to lake level, climate and different land uses in the region.*

There is a very real need to provide economic opportunity in the region, she says, particularly in the north, where poverty levels are high. But she questions the sustainability of recent initiatives to reignite forestry expansion amid the alarming Sibaya water decline.

Over time, working with several collaborators including the Department of Water and Sanitation, commercial forestry and scientists, SAEON expanded its activities into the northern section of the coastal plain and from 2014 onwards, and has also provided logistical support to a Water Research Commission (WRC) project aimed at assessing the hydrodynamics of the Sodwana Bay system.

In 2015 node staff started working in the Vazi pans area, facilitated through a WRC project run by SAEON's Prof Colin Everson, which focuses on understanding alternative agroforestry systems and plant water use. "In May 2017, I called an informal meeting of invited experts in groundwater modelling who had insight into the Sodwana-Sibayi-Vazi-Kosi systems, including experts from the Department of Water and Sanitation," she notes. "The intention was to determine if there was consensus of what might be happening in the system and if there was a case for solid long-term observation by SAEON in the area."

But because of the extended drought and water loss, problems emerged in gathering reliable data after several monitoring stations and the only lake water level gauging station became

stranded above the receding water line. SAEON has stepped in with temporary monitoring systems to help ensure a continued record.

Van Rensburg notes that poverty levels around Sibaya are amongst the highest in the country and because the soils are poor for agriculture, there appear to be limited options for economic growth.

"It is therefore imperative when initiating work in the area to ensure there is community buy-in at the outset. The socio-economic context and concern regarding the potential impacts of alternative land uses such as forestry, which many see as the only major source of income, in my view necessitates a multi-disciplinary collaborative approach if we are to see traction in response to our work."

To this end, SAEON and the Isibusiso Esihle Science Discovery Centre (a home-grown science centre close to Vazi pans), hosted an informal workshop inviting relevant stakeholders and interested experts from different disciplines including hydrology, sustainability, resources economics and horticulture.

The workshop began by listening to the voices of community members and industry operating in the area and there was unanimous concern regarding the decline in the water table and the role that plantations may be playing in this in conjunction with the drought.

"The main request was for scientists to work together with industry and community members to understand trends, but more importantly, to provide guidance on alternatives: 'We know there is a problem, but we cannot remove the trees without alternatives being in place. Please work with us to find alternatives,' was the message that emerged."

Apart from the lake itself, impacts have also radiated out into the surrounding wetland areas, including dried-out pans where crocodiles, fish and hippo were once common.

"Changes in groundwater dynamics at a regional scale will consequently have impacts for both the wetland and the terrestrial systems. The vegetation is a rich mosaic of different grassland, forest, savanna and thicket communities and hosts a high number of endemics," she explains.

Large areas of unique peatlands have also dried out, with extensive fires adding to the damage. The peatlands in this section of Maputaland, estimated to be around 7 000 years old, play a vital role in retaining moisture from heavy rainfall and releasing it gradually. And as these unique, natural sponges and wetlands dry out, Van Rensburg says it becomes more difficult to map and accurately delineate buffer zones and land which should be off-limits to timber expansion.

While the National Water Act specifically details streamflow reduction activities, she believes these provisions do not cater adequately for impacts on groundwater-driven systems such as Sibaya. Ideally, she says, policies should be revised to take account of the unique hydro-ecological nature of coastal systems and climate change viability.

Sue van Rensburg



Student, Josephine Magolego, and Sue van Rensburg of SAEON collecting water temperature samples from Lake Sibaya.



Sue van Rensburg



*This time-series sequence of images illustrates the rapid decline in water levels in the southern basin, Banda Banda Bay, of the lake since 2014*



Other potential solutions to land-use dilemmas, she suggests, could include greater community involvement in wildlife, marine and coastal tourism ventures. "Tourism will not make everyone rich. But it can provide a sustainable living for more people in the vicinity of the iSimangaliso Wetland Park World Heritage Site. This region has so much to offer and there is no reason why it should not become a rival to Kruger National Park."

According to the national Department of Environment, Forestry and Fisheries, the Mbazwana and Manzengwenya plantations are still classified as a State Forest but the biological assets were transferred to the Tembe, Mbila, Mabaso Development Trust by the Department of Agriculture, Forestry and Fisheries in April 2016.

The community trusts then formed Tembe Mbila Mabaso Forestry (TMM), to manage the plantations. TMM Forestry took over the management of the plantations and were currently carrying out all operations related to harvesting, silviculture and forest protection.

Responding to questions from *the Water Wheel* on what action had been since 2017 to implement the recommendation to remove at least 5km<sup>2</sup> of plantation forests from the Sibaya catchment, department spokesman Albi Modise said that TMM Forestry was well aware of the situation around Lake Sibaya.

"The plantations were established in the 1950s and the Department of Water Affairs and Sanitation are currently dealing with this issue. . . TMM is in the process of planting macadamia trees on 3 000 hectares of land as these trees use far less water than Eucalypts, hence the change in land use will be beneficial to Lake Sibaya."

He added that SAEON was also on board with TMM and had established a monitoring site on the first area planted with macadamias trees.

"They will establish the water use of the trees and try to provide an evidence-based approach to understanding trends to work out relative impacts of climate change. They will be establishing a site in new Eucalyptus plantings as well so that the water usage between the two different crops can be proved scientifically."

"TMM survives on timber sales and no grant funding is available to manage the plantations. For this reason, the company is intending to utilise all the available timber whilst managing the plantations in an environmentally sustainable manner."

Modise said the plantable area in the Manzengwenya plantation was 13 000 ha and that removal of 5 km<sup>2</sup> of timber would make this unit uneconomical for a timber plantation. TMM was also exploring alternative water-efficient crops, along with cattle farming and tourism.

Van Rensburg emphasises that Maputaland is a unique area requiring unique treatment and that part of the solution will depend on building a common vision, unlocking the region's ecotourism potential, combined with innovative, profitable, and climate-smart agricultural systems.

"We are moving into a new era where old rules, past trends and methodologies may no longer be appropriate. Exploring solutions as a collective with all parties represented and using social learning processes may be an important approach for ensuring long term success."



Sue van Rensburg

*Banda Banda Bay, southern basin. Mlu Shabalala, UKZN PhD student on his first trip to the region gains an understanding of how important his work is to provide evidence-based decision support for water-wise land use for the region.*



Sue van Rensburg

*High and dry. A Department of Water and Sanitation gauging station at Banda Banda bay is now stranded above the receding water level.*